Fano-Feshbach resonances observation in ultracold thulium gas

D. A. Pershin^{1,2}, V. V. Tsyganok^{1,2}, V. A. Khlebnikov¹, E. T. Davletov^{1,2,4}, I. S. Cojocaru^{1,5}, A. V. Akimov^{1,3,5}

¹Russian Quantum Center, Business Center "Ural", 100A Novaya Str., Skolkovo, Moscow, 143025, Russia ²Moscow Institute of Physics and Technology, Institutskii per. 9, Dolgoprudny, Moscow Region 141701, Russia ³PN Lebedev Institute RAS, Leninsky Prospekt 53, Moscow, 119991, Russia

⁴Center for Energy Systems, Skolkovo Institute of Science and Technology, 3 Nobel Street, Skolkovo, Moscow Region 143026, Russia

⁵Texas A&M University, 4242 TAMU, College Station, Texas, 77843, USA, e-mail: daniil.a.pershin@phystech.edu

Ultracold atomic gases are a powerful tool for quantum simulations. One of the most desirable feature of such systems is a tunable interaction between particles. Such control can be provided with scattering length variation via magnetically tuned Fano-Feshbach resonances [1]. Dipolar atoms such thulium are good candidates for quantum simulations since they have low field Fano-Feshbach resonances and a wide range anisotropic interaction [2].

We report about the s- and d-wave thulium Fano-Feshbach resonances observation in the magnetic field range between 0 and 8 Gauss at temperatures of atomic cloud 1.5 and 12 μ K. Atom ensembles were cooled and polarized with a narrow line far detuned magneto-optical trap. Further cooling and experiments occur in the optical dipole trap operating on 532 nm. The measurement of Fano-Feshbach resonances was conducted by detecting losses of atoms from the optical dipole trap, caused by a significant increase of 3-body recombination in Fano-Feshbach resonances. The resonances measured were used to determine the sign on the background scattering length of the thulium. This was done by detecting temperature anomaly near one of the strong resonances, which was associated with the zero of the scattering length [3].

Keywords: FESHBACH RESONANCES, DIPOLAR ATOMS, ULTRACOLD GASES

References

- [1] C. Chin, R. Grimm, P. Julienne, and E. Tiesinga, Rev. Mod. Phys. 82, 1225 (2010).
- [2] S. Kotochigova, Rep. Prog. Phys. 77 (2014).
- [3] K. Aikawa, A. Frisch, M. Mark, S. Baier, A. Rietzler, R. Grimm and F. Ferlaino, PRL 108, 210401 (2012).